

The role of skin biopsy in the detection of exposure to endocrine disrupting chemicals in Mediterranean cetaceans

M. CRISTINA FOSSI*, LETIZIA MARSILI* AND GIUSEPPE NOTARBARTOLO DI SCIARA[†]

Contact e-mail: cdfos@tin.it

ABSTRACT

Use of skin biopsy is proposed as a sensitive non-lethal technique for the hazard assessment of Mediterranean cetaceans exposed to endocrine disrupting chemicals (EDCs). EDCs are a structurally diverse group of compounds that may adversely affect the health of humans and wildlife or their progeny, by interaction with the endocrine system. In the Mediterranean environment top predators accumulate high concentrations of polyhalogenated aromatic hydrocarbons (PHAHs) and toxic metals, incurring high toxicological hazard. In this paper, the hypothesis that Mediterranean cetaceans are potentially at risk due to PHAH-EDCs is investigated using skin biopsy samples. Benzo- α -pyrene monooxygenase (BPMO) activity in skin biopsies was used as a potential indicator of exposure to different organochlorines (OCs) known to have endocrine disrupting properties. The main objective of this paper was to use this non-destructive ecotoxicological tool to define the potential hazard to Mediterranean odontocete and mysticete species, comparing the present data with values detected in other cetaceans from heavily polluted areas, affected by pseudohermaphroditism and other reproductive dysfunction. Subcutaneous tissue consisting of skin and blubber was obtained from striped dolphins (*Stenella coeruleoalba*), bottlenose dolphins (*Tursiops truncatus*), common dolphins (*Delphinus delphis*) and fin whales (*Balaenoptera physalus*) in the Mediterranean basin. Sampling was performed in the western Ligurian Sea, between Corsica and the French-Italian coast, and in the Ionian Sea. High concentrations of DDT metabolites and PCB congeners (known as Endocrine Disruptors) were detected in the different species. Significant differences in BPMO induction and OC levels were found between odontocetes and mysticetes. Differences in organochlorine bioaccumulation and consequently potential risk due to endocrine disruptors were primarily related to different positions in the marine food web. A statistical correlation was found between BPMO activity and organochlorine (op'DDT, a potent estrogen and antiandrogen and pp'DDE, a potent antiandrogen) levels in skin biopsy specimens of the endangered Mediterranean population of common dolphin. Several conclusions on the potential risk to Mediterranean cetaceans can be drawn from comparison of the levels of OC-EDs detected in Mediterranean odontocetes with those in white whales (*Delphinapterus leucas*) of the St Lawrence estuary and bowhead whales (*Balaena mysticetus*) affected by pseudohermaphroditism and other reproductive dysfunction. Finally, these results suggest that BPMO induction may be an early sign of exposure to EDCs such as OCs and a warning of the possibility of transgenerational effects through exposure of future generations via the placenta and milk.

KEYWORDS: BIOPSY SAMPLING; POLLUTANTS; ORGANOCHLORINES; STRIPED DOLPHIN; BOTTLENOSE DOLPHIN; COMMON DOLPHIN; FIN WHALE

INTRODUCTION

Endocrine Disrupting Chemicals (EDCs) have recently attracted much public and scientific attention (Colborn *et al.*, 1993; 1996; 1998). EDCs are a structurally diverse group of compounds that may adversely affect the health of humans and wildlife, or their progeny, by interaction with the endocrine system (Gillesby and Zacharewski, 1998). They include chemicals used heavily in the past in industry and agriculture, such as polychlorinated biphenyls and organochlorine pesticides, and also plasticisers and surfactants. Many known EDCs are estrogenic and affect reproductive function. Most xenobiotic estrogens and their metabolites are lipophilic and persistent which is why they bioaccumulate and biomagnify (Arukke *et al.*, 1997; Colborn *et al.*, 1998; Reijnders *et al.*, 1999).

Man-made EDCs range across all continents and oceans; some geographic areas are potentially more threatened than others: one of these is the Mediterranean Sea. This basin has limited exchange of water with the Atlantic Ocean, and is surrounded by some of the most heavily populated and industrialised countries in the world. Levels of some xenobiotics are much higher here than in other seas and oceans (Bernard, 1978; Ankley *et al.*, 1998). Mediterranean marine fauna could therefore be a target of EDCs. In this environment, top predators (such as large pelagic fish and marine mammals) tend to accumulate high quantities of polyhalogenated aromatic hydrocarbons (PHAHs) and toxic

metals (Leonzio *et al.*, 1992; Corsolini *et al.*, 1995; Marsili and Focardi, 1996; 1997). For example, levels of PHAHs in a top predator of the Mediterranean, the striped dolphin, are 1-2 orders of magnitude higher than in Atlantic and Pacific dolphins of the same species (Marsili, 2000). This suggests that Mediterranean top predator species, and particularly odontocetes, are potentially 'at risk' due to EDC contamination.

The potential effects of EDCs on a Mediterranean fish species of commercial interest, the swordfish (*Xiphias gladius*), were recently investigated by Fossi *et al.* (2001) using vitellogenin (Vtg) and zona radiata proteins (Zrp) as diagnostic and prognostic biomarkers. Significant induction of typically female proteins (Vtg and Zrp) was detected by ELISA and Western Blot in adult males of the species. Vtg and Zrp were found to be significantly more induced in adult male Mediterranean swordfish than 25 Atlantic specimens. These results are the first warning of the potential risk for reproductive function of Mediterranean top predators, and suggest the need for continuous monitoring of this marine environment and other Mediterranean top predator species such as marine mammals.

It is more difficult to explore the ecotoxicological risk to free-ranging Mediterranean cetaceans than to swordfish. When the levels of organochlorines in swordfish (average DDTs in blubber 2.13 $\mu\text{g/g}$ f.w.; average PCBs in blubber 1.97 $\mu\text{g/g}$ f.w.; Ausili, pers. comm.) are compared with those found in free-ranging striped dolphins, levels are found to be

* Department of Environmental Science, University of Siena, Via Mattioli 4, 53100 Siena, Italy.

[†] ICRAM, Via di Casalotti 300, 00166 Roma, Italy.

10 to 20 times higher in the latter (average DDTs in blubber 26.65 µg/g d.w.; average PCBs in blubber 46.67 µg/g d.w.; Marsili, 2000). The use of skin biopsies to assess benzo- α -pyrene monooxygenase (BPMO) activity and organochlorine body loads in cetaceans has been proposed and validated in several papers (Fossi *et al.*, 1992; 1997; 1999; 2000; Fossi and Marsili, 1997; Marsili *et al.*, 1998). The aim of this paper is to define 'potential hazard' to Mediterranean odontocete and mysticete species, using skin biopsies as a non-lethal ecotoxicological tool for the evaluation of BPMO (MFO) activity, as a potential indicator of exposure to different organochlorines (OCs) known to have endocrine disrupting properties. The final assessment involves comparing the data of Mediterranean species with OC-ED levels detected in other cetaceans from heavily polluted areas, affected by pseudohermaphroditism and other reproductive dysfunction.

MATERIALS AND METHODS

Subcutaneous tissue consisting of skin and blubber was obtained from striped dolphins, bottlenose dolphins, common dolphins and fin whales, in the western Ligurian Sea (1992-1999), between Corsica and the French-Italian coast, and in the Ionian Sea. Samples were collected from fin whales using biopsy darts launched with a 150-pound *Barnett Wildcat II* test crossbow. Dolphins were sampled while riding the vessel's bow wave by means of a pole with a biopsy tip. To avoid the possibility of infection, the biopsy tips were sterilised with ethanol before use. Biopsy specimens were taken from the dorsal area surrounding the dorsal fin. The samples were immediately placed in liquid nitrogen. The small size of the biopsy samples (between 0.200g and 0.002g) did not permit isolation of the microsomal fraction. BPMO activity was detected in the whole tissue. Since the connective tissue was very tough, the epidermis was homogenised in 1.15% KCl buffer at pH 7.5 by thermal shock and separated by freezing in liquid N₂ and pulverising with ultrasound in a Potter apparatus. BPMO activity was assessed using the incubation mixture proposed by Fossi *et al.* (1992). Each sample and the blanks were incubated in a shaking bath for 1 hour at 37°C. Activity was expressed in arbitrary units of fluorescence (A.U.F./h/g tissue). Samples of subcutaneous blubber (about 0.3g) were freeze-dried and extracted with n-hexane in a Soxhlet apparatus for analysis of chlorinated hydrocarbons. Sample purification was carried out by adding concentrated sulphuric acid to the extracts; after elimination of 'black' residues, the extracts were reconcentrated and purified by Florisil column chromatography. The analytical method used was high resolution capillary gas chromatography with a Perkin-Elmer Series 8700 GC and a 63Ni ECD. Capillary gas chromatography revealed op'- and pp'- isomers of DDT and its derivatives DDD and DDE, and about 30 PCB congeners.

The data were processed by summary statistics and ANOVA using Statgraphics software (Statistical Graphics Corporation), Statistica and Excel (Microsoft), assigning $p < 0.05$ as significance level.

RESULTS AND DISCUSSION

There are several types of organochlorine endocrine disruptors (Adami *et al.*, 1995; Kelce *et al.*, 1995; Vonier *et al.*, 1996; Wong and Pessah, 1996; Hansen, 1998; Sohoni and Sumpter, 1998; Hilscherova *et al.*, 2000) commonly found in Mediterranean cetaceans (Aguilar and Borrell,

1994; Aguilar, 2000; Marsili, 2000) (Table 1): environmental estrogens, anti-estrogens, environmental androgens and anti-androgens. Endocrine disruptors act by mimicking steroid sex-hormones, both estrogens and androgens, by binding to hormone receptors or influencing cell pathways (environmental estrogens and androgens), they also interfere with steroidogenesis, metabolism, and excretion of hormones, or by blocking and altering hormonal binding to hormone receptors (anti-estrogens, anti-androgens). Environmental estrogens have been more extensively studied than the others and may be the most common EDCs (Colborn *et al.*, 1993; 1996; 1998; Fournier *et al.*, 2000; Colborn, 2002).

Table 1

DDT metabolites and PCB congeners (known as Endocrine Disruptors) most commonly found in Mediterranean cetaceans.

| Activity | | Potency ¹ | aER binding IC50 (M) ² | ER binding IC50 (M) ³ | ER binding RBA,% ⁴ |
|---------------|----------------------------------|----------------------|-----------------------------------|----------------------------------|-------------------------------|
| DDTs | Estrogen | | | | |
| pp'DDT | Estrogen | | | | |
| | Antiandrogen | | | | |
| | ER agonist | | >50 ^a | >1000 | |
| op'DDT | Estrogen | ++ | | | |
| | Antiestrogen | + | | | |
| | Antiandrogen | +++ | | | |
| | ER agonist | | 9.1 | 5 | 0.1 |
| pp'DDE | Estrogen | + | | | |
| | Antiestrogen | + | | | |
| | Androgen | + | | | |
| | Antiandrogen | ++ | | | |
| | ER agonist | | >50 ^a | >1000 | |
| | AR agonist | | | | |
| | AR antagonist | | | | |
| op'DDE | Estrogen | | | | |
| | ER agonist | | 37.25 | | |
| pp'DDD | ER agonist | | | >1000 | |
| op'DDD | ER agonist | | 2.26 | | |
| PCBs | | | | | |
| Arochlor 1260 | Estrogen | | | | |
| | Effect on sexual differentiation | | | | |
| | Gonadal abnormalities | | | | |
| 95 | Estrogen | + | | | |
| 99 | Estrogen | ++ | | | |
| 101 | Estrogen | | | | <0.001 |
| 118 | Antiestrogen | ++ | | | |
| 153 | Estrogen | +++ | | | 0.004 |

¹The most potent chemical for each activity was assigned a potency of four plus signs (++++), and the potency of all the chemicals was expressed relative to this (Hansen, 1998; Sohoni & Sumpter, 1998).

²Inhibitor concentrations necessary for 50% inhibition (IC50) of 3H17-estradiol binding to estrogen receptors (aER) in alligators. The aER binding IC50 value for 17-estradiol was 0.0078M.

³Compounds that inhibited 3H17-estradiol but were insoluble at concentrations necessary to achieve 50% inhibition (Vonier *et al.*, 1996).

⁴Inhibitor concentrations necessary for 50% inhibition (IC50) of 3H17-estradiol binding to ERs in rats. The ER binding IC50 value for 17-estradiol was 0.002M (Kelce *et al.*, 1995).

⁵Relative estrogen receptor-binding affinities (RBA). Competitive binding with estradiol in rat uterine ER preparations (Hansen, 1998).

Activity references: Adami *et al.*, 1995; Wong & Pessah, 1996; Hansen, 1998; Sohoni & Sumpter, 1998; Hilscherova *et al.*, 2000).

High concentrations of organochlorines with known EDC properties (Table 1) were detected in the different cetaceans (Table 2). Significant differences (T-student test = $p < 0.05$) in levels were found between odontocetes and mysticetes.

Table 2

Arithmetic mean (ng/g fresh weight) and standard deviations (in brackets) of DDT metabolites and PCB congeners known as EDCs in the four species of Mediterranean cetaceans.

| | Fin whale (n=63) | Striped dolphin (n=95) | Bottlenose dolphin (n=7) | Common dolphin (n=13) |
|-------------|----------------------------------|------------------------------------|------------------------------------|------------------------------------|
| pp'DDT | 457.2 (301.2) | 2676.9 (2058.3) | 793.2 (892.8) | 336.4 (230.2) |
| op'DDT | 438.9 (348.5) | 1306.1 (1085.9) | 450.2 (322.1) | 400.1 (297.5) |
| pp'DDE | 3679.4 (2560.3) | 19972.2 (17557.2) | 10138.5 (9973.8) | 13453.6 (11437.1) |
| op'DDE | 45.4 (27.6) | 288.4 (277.5) | 286.2 (227.5) | 99.6 (95.8) |
| pp'DDD | 446.2 (318.4) | 1884.1 (1472.9) | 543.9 (381.3) | 467.1 (376.7) |
| op'DDD | 101.5 (64.1) | 521.7 (604.6) | 71.5 (104.0) | 115.5 (205.8) |
| DDTs | 5168.6 (3844.4) | 26649.4 (21391.8) | 12283.5 (10532.3) | 14872.3 (11801.4) |
| PCB 95 | 154.9 (73.1) | 533.8 (343.4) | 566.5 (402.6) | 344.4 (389.2) |
| PCB 101 | 144.8 (70.4) | 579.3 (381.2) | 562.5 (449.5) | 219.9 (179.8) |
| PCB 99 | 372.4 (275.8) | 1879.2 (1668.5) | 820.6 (759.8) | 560.2 (313.1) |
| PCB 118 | 203.9 (102.8) | 994.9 (594.4) | 951.0 (718.2) | 1788.1 (1710.9) |
| PCB 153 | 1100.8 (529.6) | 8592.5 (5174.3) | 8186.7 (9376.4) | 4214.7 (2072.1) |
| PCBs | 6597.4 (3270.8) | 49127.6 (29669.2) | 31446.5 (25129.7) | 20025.8 (9024.3) |

Highest levels were found in striped dolphins, followed by bottlenose dolphins and common dolphins. Organochlorine values were lower in fin whales than in the dolphin species by a factor of 2-8. Differences in organochlorine bioaccumulation and consequently potential risk due to endocrine disruptors are primarily related to different positions in the marine food web. In this case, the odontocete species may be regarded as terminal consumers that prey mainly on fish and squid; fin whales occupy a lower level, feeding on plankton which contain average PCB levels 5-10 times lower than odontocete prey. This suggests that odontocetes incur greater risk than Mediterranean mysticetes.

Levels of the different OCs with endocrine disrupting properties need to be explored separately (Table 1). The high percentage of op'DDT detected in fin whale samples (Fig. 1a-b) is noteworthy, because this DDT metabolite is a potent oestrogen and anti-androgen and could theoretically affect the low reproductive rate of this mysticete.

This research was prompted by the fact that some species, such as the common dolphin, have almost completely disappeared from the Mediterranean Sea. In considering the potential use of this ecotoxicological tool (skin biopsy) to study inter-species susceptibility to contaminants in Mediterranean cetaceans, a high statistical correlation (Pearson product-moment correlations $p < 0.05$) was found between BPMO activity and organochlorine levels (DDTs) in skin biopsy specimens of the endangered Mediterranean population of common dolphin. op'DDT is a potent Oestrogen and anti-androgen (Table 1) (Fig. 2a) and pp'DDE is a potent anti-androgen (Table 1) (Fig. 2b). This data suggests that EDCs may be one stress factor for common dolphin populations in the Mediterranean Sea.

In an attempt to define the 'potential hazard' of Mediterranean odontocete and mysticete species, a preliminary assessment can be undertaken, comparing the data of the Mediterranean species with those for PCB-EDS detected in other cetaceans from heavily polluted areas, affected by pseudohermaphroditism and other reproductive dysfunction.

Research in other seas and oceans indicates that some cetaceans, particularly odontocetes, have detectable and sometimes extremely high levels of substances known or suspected to be EDCs, such as PCBs, DDTs, chlorinated pesticides, brominated flame retardants and tributyltin (TBT). Biologists have reported a range of effects of these chemicals on cetaceans, including immunosuppression, cancer, skin lesions, secondary infections and diseases, sporadic die-offs, and reduced reproductive success. There are several examples suggesting that exposure to OC insecticides and PCBs has affected endocrine function and reproduction in marine mammals. For example, transformation of epididymal and testicular tissue has been observed in North Pacific common minke whales, *Balaenoptera acutorostrata* (Fujise *et al.*, 1998).

Another example is the endangered white whales of the St Lawrence estuary, now among the most contaminated animals on earth, with tumours and reproductive problems




Fig. 1a. Percentage of DDT compounds known to be EDCs (see Table 1) out of total DDTs in skin biopsy specimens from Mediterranean cetaceans.




Fig. 1b. Percentage of PCB congeners known to be EDCs (see Table 1) out of total PCBs in skin biopsy specimens from Mediterranean cetaceans.

(Deguise *et al.*, 1995; Martineau *et al.*, 1999; 2002). De Guise *et al.* (1994) reported a true hermaphrodite white whale. This animal had two testicles, two separate ovaries and the complete ducts of each sex; cervix, vagina and vulva were absent. Mature spermatozoa were found in the lumen of seminiferous tubules in the testicles, and many involuted corpora lutea were recognised in the ovaries.

Increasing residue levels of PCBs and DDE in the blubber of Dall's porpoises (*Phocoenoides dalli*) were found to have a negative effect on testosterone levels in blood. Testosterone levels decreased in a statistically significant way with increasing DDE concentrations. These results suggest that current levels of environmental contamination by persistent organochlorines can cause an imbalance in sex hormones and subsequent reproductive abnormalities in wild





Fig. 2. Statistical correlation ($p < 0.05$) between BPMD activity and organochlorine levels: (A) (op'DDT) and (B) (pp'DDE), in common dolphin skin biopsies.

porpoises. The other hormone measured, aldosterone, which has no sexual function, showed levels which were independent of the effects of PCBs and DDE (Subramanian *et al.*, 1987).

In the endangered bowhead whale, not all sexual activity leads to conception. Pseudohermaphroditism has been reported in at least two males with testicular feminisation (Tarpley *et al.*, 1995). This is a relatively high incidence, as only 76 bowheads were closely examined between 1980 and 1989 (Philo *et al.*, 1993). Concentrations of most organochlorine contaminants in biological material of 20 bowhead whales from Barrow (Alaska, USA) were low compared to those in tissues of other cetaceans, especially odontocetes, but it is impossible to rule out any link between pseudohermaphroditism and these contaminants.

Despite difficulties in establishing direct cause-effect links (Reijnders *et al.*, 1999) it is worth noting that the levels of PCBs found in Mediterranean odontocetes are similar to those detected in hermaphrodite white whales of the St Lawrence estuary (Fig. 3a); levels of PCBs detected in Mediterranean fin whales are approximately 10 times higher




Fig. 3a. Comparison of PCB levels in Mediterranean odontocetes (striped, bottlenose and common dolphins) and odontocete species affected by hermaphroditism. Broken line — levels of PCBs found in populations of white whale manifesting hermaphroditism.




Fig. 3b. Comparison of PCB levels in a Mediterranean fin whale and mysticete species manifesting pseudohermaphroditism. Broken line — levels of PCBs found in populations of bowhead whale affected by pseudohermaphroditism.

than those found in bowhead whales with pseudohermaphroditism and other reproductive dysfunctions (Fig. 3b). This observation illustrates the potential risk to which these species are exposed in the Mediterranean Sea. Future research on stranded animals will help to clarify potential effects of these chemicals on gonad integrity.

Inter-species differences in susceptibility to EDCs will be a major topic to consider in this context. Future studies should explore the role of detoxification enzymes and estrogen receptors (ER) in inter-species susceptibility to EDCs. As an approach to this problem, the authors have been studying cultures of fibroblasts from different cetaceans (striped dolphin, bottlenose dolphin, common dolphin and fin whale; Marsili *et al.*, 2000) sampled in the Mediterranean Sea between 1999 and 2001. Skin biopsy samples were stored in a 'cell medium' and transported to the lab within 24-36 hours where they were processed to obtain fibroblast cultures. Fibroblasts of the different species have already been cultured, and when the number of cells is sufficient for testing, inter-species differences in susceptibility to the main Mediterranean OCs with EDC properties will be investigated. The results will be a first indication of any differences in the susceptibility of the various Mediterranean cetaceans to this dangerous class of chemicals.

ACKNOWLEDGEMENTS

This project was financed by a grant from the Italian Ministry for the Environment and ICRAM. We thank the Tethys Research Institute and Giovanni Neri for scientific and technical support.

REFERENCES

- Adami, H.O., Lipworth, L., Titus-Ernstoff, L., Chung-Cheng, H., Hamberg, A., Aahborg, U., Baron, J. and Trichopoulos, D. 1995. Organochlorine compounds and estrogen-related cancers in women. *Cancer Causes and Control* 6:551-66.
- Aguilar, A. 2000. Population biology, conservation threats and status of Mediterranean striped dolphins (*Stenella coeruleoalba*). *J. Cetacean Res. Manage.* 2(1):17-26.
- Aguilar, A. and Borrell, A. 1994. Abnormally high polychlorinated biphenyl levels in striped dolphins (*Stenella coeruleoalba*) affected by the 1990-1992 Mediterranean epizootic. *Sci. Total Environ.* 154(2-3):237-47.
- Ankley, G., Mihaich, E., Stahl, R., Tillitt, D., Colborn, T., McMaster, S., Miller, R., Bantle, J., Campbell, P., Denslow, N., Dickerson, R., Folmar, L., Fry, M., Giesy, J., Gray, L.E., Guiney, P., Hutchinson, T., Kennedy, S., Kramer, V., LeBlanc, G., Mayes, M., Nimrod, A., Patino, R., Peterson, R., Purdy, R., Ringer, R., Thomas, P., Touart, L., Van Der Kraak, G. and Zacharewski, T. 1998. Overview of a workshop on screening methods for detecting potential (anti)estrogenic/androgenic chemicals in wildlife. *Environ. Toxicol. Chem.* 17(1):68-87.
- Arukke, A., Knudsen, F.R. and Goksoyr, A. 1997. Fish zona radiata (eggshell) protein: a sensitive biomarker for environmental estrogens. *Environ. Health Perspect.* 105:418-22.
- Bernard, M. 1978. *Ocean Management*. 253pp.
- Colborn, T. 2002. Clues from wildlife to create an assay for thyroid system disruption. *Environ. Health Perspect.* 110(3):363-7.
- Colborn, T., vom Saal, F.S. and Soto, A.M. 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ. Health Perspect.* 101:378-84.
- Colborn, T., Dumanoski, D. and Myers, J.P. 1996. *Our Stolen Future*. Dutton, Penguin Books USA, New York. 306pp.
- Colborn, T., Smolen, M.J. and Rolland, R. 1998. Environmental neurotoxic effects: the search for new protocols in functional teratology. *Toxicol. Ind. Health* 14(1/2):9-23.
- Corsolini, S., Focardi, S., Kannan, K., Tanabe, S., Borrell, A. and Tatsukawa, R. 1995. Congener profile and toxicity assessment of polychlorinated biphenyls in dolphins, sharks and tuna collected from Italian coastal waters. *Mar. Environ. Res.* 40(1):33-53.
- Deguisse, S., Bisailon, A., Seguin, B. and Lagace, A. 1994. The anatomy of the male genital system of the beluga whale, *Delphinapterus leucas*, with special reference to the penis. *Anatomia Histologia Embryologia - Journal of Veterinary Medicine Series C - Zentralblatt Fur Veterinarmedizin Reihe C* 23(3):207-16.
- Deguisse, S., Martineau, D., B eland, P. and Fournier, M. 1995. Possible mechanisms of action of environmental contaminants on St. Lawrence beluga whales (*Delphinapterus leucas*). *Environ. Health Perspect.* 103(4):73-7.
- Fossi, M.C. and Marsili, L. 1997. The use of non-destructive biomarkers in the study of marine mammals. *Biomarkers* 2:205-16.
- Fossi, M.C., Marsili, L., Leonzio, C., Notarbartolo-di-Sciara, G., Zanardelli, M. and Focardi, S. 1992. The use of non-destructive biomarkers in Mediterranean cetaceans - preliminary data on MFO activity in skin biopsy. *Mar. Poll. Bull.* 24(9):459-61.
- Fossi, M.C., Savelli, C., Marsili, L., Casini, S., Jimenez, B., Junin, M., Castello, H. and Lorenzani, J.A. 1997. Skin biopsy as a nondestructive tool for the toxicological assessment of endangered populations of pinnipeds: preliminary results on mixed function oxidase in *Otaria flavescens*. *Chemosphere* 35(8):1623-35.
- Fossi, M.C., Casini, S. and Marsili, L. 1999. Nondestructive biomarkers of exposure to endocrine-disrupting chemicals in endangered species of wildlife. *Chemosphere* 39(8):1273-85.
- Fossi, M.C., Marsili, L., Neri, G., Casini, S., Bearzi, G., Politi, E., Zanardelli, M. and Panigada, S. 2000. Skin biopsy of Mediterranean cetaceans for the investigation of interspecies susceptibility to xenobiotic contaminants. *Mar. Environ. Res.* 50:517-21.
- Fossi, M.C., Casini, S., Ancora, S., Moscatelli, A., Ausili, A. and Notarbartolo di Sciara, G. 2001. Do endocrine disrupting chemicals threaten Mediterranean swordfish? Preliminary results of vitellogenin and zona radiata proteins in *Xiphias gladius*. *Mar. Environ. Res.* 52:477-83.
- Fournier, M., Degas, V., Colborn, T., Omara, F.O., Denizeau, F., Potworowski, E.F. and Brousseau, P. 2000. Immunosuppression in mice fed on diets containing beluga whale blubber from the St Lawrence estuary and the Arctic populations. *Toxicology Letters* 112-113:311-7.
- Fujise, Y., Zenitani, R. and Kato, H. 1998. An examination of the W-stock hypothesis for North Pacific minke whales with special reference to some biological parameters using data collected from JARPEN surveys from 1994 to 1997. Paper SC/50/RMP12 presented to the IWC Scientific Committee, April 1998, Oman (unpublished). [Paper available from the Office of this Journal].
- Gillesby, B. and Zacharewski, T. 1998. Exoestrogens: mechanisms of action and strategies for identification and assessment. *Environ. Toxicol. Chem.* 17(1):3-14.
- Hansen, L.G. 1998. Stepping backward to improve assessment of PCB congener toxicities. *Environ. Health Perspect.* 106(1):171-89.
- Hilscherova, K., Machala, M., Kannan, K., Blankenship, A.L. and Giesy, J.P. 2000. Cell bioassays for detection of aryl hydrocarbon (AhR) and estrogen receptor (ER) mediated activity in environmental samples. *Environ. Sci. Pollut. Res. Int.* 7(3):159-71.
- Kelce, W.R., Stone, C.R., Laws, S.C. and Gray, L.E. 1995. Persistent DDT metabolite pp'DDE is a potent androgen receptor antagonist. *Nature, Lond.* 375:581-6.
- Leonzio, C., Focardi, S. and Fossi, C. 1992. Heavy metals and selenium in stranded dolphins of the northern Tyrrhenian (NW Mediterranean). *Sci. Total Environ.* 119:77-84.
- Marsili, L. 2000. Lipophilic contaminants in marine mammals: review of the results of ten years work at the Department of Environmental Biology, Siena University (Italy). The control of marine pollution: current status and future trends. *Int. J. Environ. Pollut.* 13:416-52.
- Marsili, L. and Focardi, S. 1996. Organochlorine levels in subcutaneous blubber biopsies of fin whales *Balaenoptera physalus* and striped dolphins *Stenella coeruleoalba* from the Mediterranean Sea. *Environ. Pollut.* 91(1):1-9.
- Marsili, L. and Focardi, S. 1997. Chlorinated hydrocarbon HCB, DDTs and PCBs levels in cetaceans stranded along the Italian coasts: An overview. *Environ. Monitoring and Assessment* 45(2):129-80.
- Marsili, L., Fossi, M.C., Notarbartolo di Sciara, G., Zanardelli, M., Nani, B., Panigada, S. and Focardi, S. 1998. Relationship between organochlorine contaminants and mixed function oxidase activity in skin biopsy specimens of Mediterranean fin whales (*Balaenoptera physalus*). *Chemosphere* 37(8):1501-10.
- Marsili, L., Fossi, M.C., Neri, G., Casini, S., Gardi, C., Palmeri, S., Tarquini, E. and Panigada, S. 2000. Skin biopsies for cell cultures from Mediterranean free-ranging cetaceans. *Mar. Environ. Res.* 50(1-5):523-6.
- Martineau, D., Lair, S., De Guise, S., Lipscomb, T.P. and B eland, P. 1999. Cancer in beluga whales from the St Lawrence Estuary, Quebec, Canada: a potential biomarker of environmental contamination. *J. Cetacean Res. Manage.* (special issue) 1:249-65.

- Martineau, D., Lemberger, K., Dallaire, A., Labelle, P., Lipscomb, T.P., Michel, P. and Mikaelian, I. 2002. Cancer in wildlife, a case study: beluga from the St Lawrence estuary, Quebec, Canada. *Environ. Health Perspect.* 110(3):285-92.
- Philo, L.M., Shotts, E.B. and George, J.C. 1993. Morbidity and mortality. pp. 275-312. In: J.J. Burns, J.J. Montague and C.J. Cowles (eds.) *Special Publication*. No. 2. *The Bowhead Whale*. 1st. Edn. Society for Marine Mammalogy, Lawrence, Kansas.
- Reijnders, P.J.H., Donovan, G.P., Aguilar, A. and Bjørge, A. 1999. Report of the Workshop on Chemical Pollution and Cetaceans, March 1995, Bergen, Norway. *J. Cetacean Res. Manage.* (special issue) 1:1-42.
- Sohoni, P. and Sumpter, J.P. 1998. Several environmental oestrogens are also anti-androgens. *J. Endocrinol.* 158:327-39.
- Subramanian, A.N., Tanabe, S., Tatsukawa, R., Saito, S. and Miyazaki, N. 1987. Reduction in the testosterone levels by PCBs and DDE in Dall's porpoises of the northwestern North Pacific. *Mar. Poll. Bull.* 18(12):643-6.
- Tarpley, R.J., Jarrell, G.H., George, J.C., Cabbage, J. and Stott, G.G. 1995. Male pseudohermaphroditism in the bowhead whale, *Balaena mysticetus*. *J. Mammal.* 76(4):1267-75.
- Vonier, P.M., Crain, D.A., McLachlan, J.A., Guillette, L.J., Jr. and Arnold, S.F. 1996. Interaction of environmental chemicals with the estrogen and progesterone receptor from the oviduct of the American alligator. *Environ. Health Perspect.* 104(12):1318-22.
- Wong, P.W. and Pessah, I.N. 1996. *Ortho*-substituted PCBs alter calcium regulation by a ryanodine receptor-mediated mechanism: structural specificity towards skeletal- and cardiac-type microsomal calcium release channels. *Mol. Pharmacol.* 49:740-51.

Date received: March 2002.

Date accepted: August 2002.